

**Comment on the paper by Arik, Gün and Yildiz (hep-th/0212129)
on "Invariance Quantum Group of the Fermionic Oscillator"**

R.Parthasarathy¹

and

K.S.Viswanathan²

Department of Physics, Simon Fraser University
Burnaby, B.C., Canada V5A 1S6.

Abstract

A non-trivial q-deformation of the fermionic oscillator proposed by us in 1991 is recalled in view of the result of the paper cited in the title.

In the paper by Arik, Gün and Yildiz [1], the authors considered the fermionic oscillator

$$\begin{aligned} cc^* + c^*c &= 1, \\ c^2 &= 0. \end{aligned} \tag{1}$$

and state that the above algebra (1) does not admit a q-deformation. Then, they have shown that the structure of the inhomogeneous invariance group of fermionic oscillator is a quantum group.

The fact that when $c^2 = 0$, the fermionic oscillator algebra does not admit a q-deformation has been pointed out in 1991 by Jing and Xu [2]. Considering a *possible* q-deformation of the fermion oscillator algebra proposed by Chaichian and Kullish [3], namely,

$$\begin{aligned} c_q c_q^* + q c_q^* c_q &= q^N, \\ c_q^2 &= 0, \quad [N, c_q^*] = c_q^*, \quad [N, c_q] = -c_q, \end{aligned} \tag{2}$$

¹Permanent address: The Institute of Mathematical Sciences, Chennai 600 113, India:
e-mail: sarathy@imsc.ernet.in

²e-mail address: kviswana@sfu.ca

and making a transformation $c = q^{-\frac{N}{2}} c_q$, $c^* = c_q^* q^{-\frac{N}{2}}$, the above algebra (2) reduces to (1), showing that (1) does not admit a q-deformation such as in (2).

However, we [4] in 1991, have proposed a *non-trivial* q-deformation of the fermion oscillator algebra (1), as

$$\begin{aligned} f_q f_q^\dagger + \sqrt{q} f_q^\dagger f_q &= q^{-\frac{N}{2}}, \\ [N, f_q^\dagger] &= f_q^\dagger, \quad [N, f_q] = -f_q; \quad f_q^2 \neq 0, \end{aligned} \quad (3)$$

which cannot be reduced to (1) and hence non-trivial. In fact, we have pointed out that in q-deforming the fermionic oscillator algebra, the Pauli principle has to be relaxed. *The relation $f_q^2 \neq 0$, allows more than two q-fermions in a given quantum state.* The Fock space of such q-fermion algebra has been constructed in [4] using fermionic q-numbers,

$$[n]_q^F = \frac{q^{-\frac{n}{2}} - (-1)^n q^{\frac{n}{2}}}{q^{-\frac{1}{2}} + q^{\frac{1}{2}}}, \quad (4)$$

and in the limit $q \rightarrow 1$, the Fock space reduces to vacuum and one-fermion state, thereby recovering the Pauli principle in this limit. So it is possible to q-deform the fermionic oscillator algebra (1) and such a q-deformed algebra is (3).

Further, using (7), (8) and (28) of [1], it follows that the action of their A matrix in (5), produces same algebra as in (2) (upto overall normalization) *but* with $(c')^2 \neq 0$, whose structure is different from (3).

References.

1. M.Arik, S.Gün and A.Yildiz, *Invariance quantum group of fermionic oscillator*, hep-th/02121219.
2. S.Jing and J.Xu, J.Phys.A: Math.Gen. **24** (1991) L891.
3. M.Chaichian and P.I.Kullish, CERN Preprint, Th.5969/90, November 1990.
4. R.Parthasarathy and K.S.Viswanathan, J.Phys.A: Math.Gen. **24** (1991) 613.